



## ARTÍCULO ORIGINAL

### VALUE OF ENTOMOLOGICAL COLLECTIONS IN THE STUDY OF BIODIVERSITY: EXAMPLES FROM THE COLLECTION OF HARD SCALES (HEMIPTERA: DIASPIDIDAE), WHITEFLIES (HEMIPTERA: ALEYRODIDAE) AND SWEAT BEES (HYMENOPTERA: HALICTIDAE) AT THE CALIFORNIA ACADEMY OF SCIENCES

John W Dooley, Allan H Smith-Pardo\*

\* Corresponding author:

United States Department of Agriculture  
Animal and Plant Health Inspection Service  
Plant Protection and Quarantine  
389 Oyster Point Blvd, Suite 2A  
South San Francisco, CA 94080  
allan.h.smith-pardo@aphis.usda.gov

#### **Abstract.**

This paper demonstrates the value of entomological collections in the study of biodiversity citing examples from material that was deposited at the California Academy of Sciences in San Francisco that resulted into the description of new insect species. Specimens used from this collection include whiteflies (Hemiptera-Sternorrhyncha-Aleyrodidae), armored scales (Hemiptera-Sternorrhyncha-Coccoidea-Diaspididae) and sweat bees (Hymenoptera-Halictidae).

**Key words:** Apoidea, Coccoidea, Entomological collection, *Gadigaleyrodes*, Taxonomy.

## Resumen.

En el presente artículo demostramos la importancia que poseen las colecciones de entomología para el estudio de la diversidad, en particular, se proveen ejemplos de material que estaba depositado como misceláneo en las colecciones de la Academia de Ciencias de California en San Francisco. Las partes de la colección utilizadas para los ejemplos incluyen las moscas blancas (Aleyrodidae), escamas duras (Diaspididae) y abejas del sudor (Halictidae).

**Palabras clave:** Apoidea, Coccoidea, Colecciones entomológicas, Gadigaleyrodes, taxonomía.

## Introduction:

### Work and new species found in miscellaneous material at the California Academy of Sciences

The California Academy of Sciences collection (CAS) has one of the largest entomological collections in the United States; in fact, their holdings were estimated to more than 15 million specimens (Norman Penny, pers. comm), which represents twice the number reported in 2004 by Suarez and Tsutsui. The CAS provides important resources to the study of the diversity of insects worldwide. Museums provide important functions such as receiving, curating, preserving, and identifying specimens while maintaining associated data (such as field notes, geographical coordinates, host records, and valuable biological information) for future research (Ross 1950, Winston 2007). In addition, CAS

provides other services such as research, education, and public outreach.

In general the biological collections at CAS maintain and keep biological data up to date and that document past and present-day patterns of biological distribution and diversity in our planet and are primary source for taxonomic work (such as taxonomic revisions and species description and name holding) as well as other biological disciplines.

While working in the curation of some sections of the CAS the authors discovered several undescribed species of armored scales (Hemiptera-Sternorrhyncha-Coccoidea- Diaspididae), whiteflies (Hemiptera-Sternorrhyncha-Aleyrodidae) and sweat bees (Hymenoptera- Halictidae) some of which have been described and published as new to science in recent years. The armored scale *Protomorgania koebelei* Dooley and Evans (2012) (figure 1) and the whitefly *Gadigaleyrodes froggatti* Dooley and Gillespie (2013) (figure 2) were recently described even though they were collected more than a hundred years ago, This contribution aims to highlight the importance of natural history collections, and in particular that of the CAS, as holders of uncounted new species awaiting for description.

### Origins of material and the process of discovery

The specimens referred herein are deposited at the CAS (Norman Penny, Collection Manager) and which represent examples of the entomological fauna for every zoogeographical region of the world.

At the beginning of this work, miscellaneous armored scales and whiteflies were slide mounted from specimens with their associated hosts using the techniques presented by Dooley (2011) since the material for this two families were found only dried without slide mounted specimens, including material collected by Froggatt (Froggatt 1932) and Koebele more than a century ago.

Another source of armored scales for some of the remarkable species found (and slide mounted) came from fresh material collected during 2002 as part of the Madagascar Diversity Project directed by the California Academy of Sciences (CAS) by Brian Fisher (PI). These specimens were preserved in 95% alcohol and sent to the University of Massachusetts at Amherst for non-destructive DNA extraction and sequencing. Based on a combination of information from two nuclear loci (28S, EF1a), one mitochondrial locus (COI-II), and morphological data, two new species each of genus *Melissoaspis* and *Melanaspis* have been described and are being published (S. Schneider pers. comm.).

In the case of the bees, just to cite an example one of us (AHSP) discovered a new species of bee when sorting out undetermined Halictidae bees; *Habralictus insularis* Smith-Pardo, 2009 (figure 3) described and some interesting remarks about its biogeography were also published by Smith-Pardo (2005) among some remarkable things, this was the first species of the genus *Habralictus* published in more than 30 years, and represented the second known species in the same genus inhabiting an island.

## Conclusions

Natural history collections, such as the one at CAS are important not only as holder of material for morphological or taxonomical studies, but also as a source of material for molecular studies that lead to the development of molecular diagnostic tools for stages of some groups of insects, that cannot be identified morphologically to species, such as adult whiteflies or immature armored scales.

We believe that by offering the examples above we emphasize even more the enormous importance of natural history collections and their care, continually curating and studying collections including digitization of data and imaging identified taxa, revising lists of established taxa with newly described genera and species, and better documentation of geographical distribution and host preferences of all species are a most of we want to keep a collection active and “alive”.

## Acknowledgments

We are grateful to Dr. Norman Penny, collection manager of the entomological collection at CAS for allowing us into the collection and for all the logistics. We will also like to thank Dr. Kyle Beucke USDA- APHIS- PPQ, Ray Gill (California Department of Food and Agriculture) for reviewing this paper, and Mr. Arthur Berlowitz, officer in charge at the Plant Inspection Station in South San Francisco for his continuous support.

## Literature Cited

Dooley J. 2011. Whitefly pupa of the

World. Plant Protection and Quarantine, United States Department of Agriculture, Available at <http://keys.lucidcentral.org/keys/v3/whitefly/Default.htm> (accessed 6 May 2012).

Dooley JW, Evans GA. 2012. A new genus and species of armored scale insect (Hemiptera: Diaspididae) from Australia found in the historic Koebele Collection of the California Academy of Sciences. *Insecta Mundi* 0218: 1-14.

Dooley JW, Gillespie PS. 2013. *Gadigaleyrodes froggatti*, a new genus and species of whitefly (Hemiptera: Aleyrodidae) from Australia. *Zootaxa* 3608 (2): 137-146.

Froggatt WW. 1932. The curators and botanists of the Botanic Gardens, Sydney. *Royal Australian Historical Society* xviii: 101-133.

Ross ES. 1950. The role of the entomological Museum. *The Pan-Pacific Entomologist* 26 (1): 1-10.

Smith-Pardo AH. 2009. A new species of *Habralictus* (Hymenoptera, Halictidae) from the Island of Grenada (Lesser Antilles) with comments on the insular species of the genus. *Zookeys* 27: 51-58.

Suarez AV, Tsutsui ND. 2004. The value of museum collections for research and society. *BioScience* 54 (1): 67-74.

Winston JE. 2007. Archives of a small planet: The significance of museum collections and museum-based research in invertebrate taxonomy. *Zootaxa* 1668: 47-54.

**Table 1.** List of whiteflies and hard scales at the CAS: first genus (two species) corresponds to Aleyrodidae, all other genera are Diaspididae. Generic names in bold letters have more than one species.

Genus	Species	Author	Host	Origin
<i>Melissoaspis</i>	<i>fisheri</i>	Ben-Dov	undetermined	Madagascar
	<i>reticulata</i>	Ben-Dov	undetermined	Madagascar
<i>Melanaspis</i>	<i>madagascariensis</i>	(Mamet)	undetermined	Madagascar
<i>Morganella</i>	<i>formicaria</i>	Ben-Dov	undetermined	Madagascar
<i>Gadigaleyrodes</i>	<i>froggatti</i>	Dooley & Gillespie	“Syncarpia”	Australia
<i>Acutaspis</i>	<i>albopicta</i>	(Cockerell)	<i>Rosa</i> sp.	Mexico
<i>Adiscoiorinia</i>	<i>secreta</i>	Green	<i>Nephelium</i> sp.	Hong Kong
<i>Andaspis</i>	sp. ( <i>A. crawii</i> ?)	NA	<i>Quercus cuspidata</i>	Japan
<i>Aonidia</i>	sp.	NA	<i>Quercus salicifolia</i>	Hong Kong
<i>Aonidiella</i>	<i>aurantii</i>	(Maskell)	<i>Olea europaea</i>	Australia
	<i>citrina</i>	(Coquillet)	<i>Cydonia</i> sp.	Western Samoa
	<i>orientalis</i>	Newstead	<i>Erythrina variegata</i>	Hong Kong
<i>Aulacaspis</i>	<i>vitis</i>	(Green)	undetermined	Hong Kong
<i>Aulacaspis</i>	<i>vitis</i>	(Green)	<i>Eleagnus</i> sp.	Taiwan
<i>Chortinaspis</i>	<i>bilobis</i>	(Maskell)	grass	Hong Kong
<i>Chrysomphalus</i>	<i>aonidum</i>	Linnaeus	undetermined	Taiwan
	<i>dictyospermi</i>	(Morgan)	<i>Citrus</i> sp.	Taiwan
<i>Diaspidiotus</i>	<i>liquidambaris</i> ?		<i>Petrea</i> sp.	Mexico
	<i>perniciosus</i>	(Comstock)	<i>Pyrus achras</i>	Hong Kong
<i>Duplaspidiotus</i>	<i>claviger</i>	(Cockerell)	<i>Hakea</i> sp.	Australia?
<i>Epidiaspis</i>	<i>leperii</i>	((Signoret)	undetermined	California
<i>Epidiaspis</i>	<i>leperii</i>	(Signoret)	<i>Malus</i> sp.	California
<i>Fiorinia</i>	sp.	NA	<i>Juniperus</i> sp.	Taiwan
	sp.	NA	undetermined	undetermined
	sp.	NA	<i>Torreya nucifera</i>	Japan
<i>Hemiberlesia</i>	<i>lataniae</i>	(Signoret)	<i>Celtis occidentalis</i>	Mexico
	<i>lataniae</i>	(Signoret)	<i>Opuntia</i> sp.	Mexico
	<i>lataniae</i>	(Signoret)	undetermined	Mexico
	<i>lataniae</i>	(Signoret)	<i>Nicotiana</i> sp.	Mexico
	<i>lataniae</i>	(Signoret)	<i>Phoradendron</i> sp.	Mexico
	<i>lataniae</i>	(Signoret)	undetermined	Mexico
	<i>lataniae</i>	(Signoret)	“Copal”	Mexico
	<i>lataniae</i>	(Signoret)	undetermined	Fiji
	<i>rapax</i>	(Comstock)	<i>Cereus</i> sp.	Mexico


	<i>rapax</i>	(Comstock)	<i>Asclepias</i> sp.	Mexico
	<i>rapax</i>	(Comstock)	<i>Mimosa</i> sp.	Mexico
	<i>rapax</i>	(Comstock)	undetermined	Mexico
	<i>rapax</i>	(Comstock)	undetermined	undetermined
	<i>rapax</i>	(Comstock)	<i>Malus</i> sp.	California
	<i>rapax</i>	(Comstock)	<i>Hedera</i> sp.	California
	<i>rapax</i>	(Comstock)	<i>Arctostaphylos</i> sp.	California
<i>Hemiberlesia</i>	<i>rapax</i>	(Comstock)	undetermined	undetermined
<i>Lepidosaphes</i>	<i>beckii</i>	(Newman)	<i>Myrtus</i> sp.	California
	<i>beckii</i>	(Newman)	<i>Citrus sinensis</i>	Sri Lanka
	<i>conchiformis</i>	(Gmelin)	<i>Crataegus</i> sp.	Mexico
	<i>mexicana</i>	(Cockerell)	Nettle tree	Mexico
	<i>newsteadi</i>	(Sulc)	<i>Sciadopitys verticillata</i>	Japan
	sp.	NA	<i>Panicum</i> sp.	Hong Kong
	<i>tubulorum</i>	Ferris	<i>Melia</i> sp.	China
	<i>ulmi</i>	(Linnaeus)	<i>Myrtus</i> sp.	California
<i>Lopholeucaspis</i>	<i>japonica</i>	Cockerell	<i>Celtis occidentalis</i>	Hong Kong
<i>Melanaspis</i>	<i>obscura</i>	(Comstock)	<i>Acer</i> sp.	Mexico
	<i>smilacris</i>	(Cockerell)	undetermined	New York
	sp.	NA	<i>Quercus</i> sp.	Arizona
	sp.	NA	<i>Syringa</i> sp.	Mexico
<i>Mycetaspis</i>	<i>personata</i>	(Comstock)	<i>Prosopis</i> sp.	Mexico
<i>Oceanaspidiotus</i>	<i>spinus</i>	(Comstock)	undetermined	Mexico
<i>Odonaspis</i>	<i>oshimaensis</i>	Kuwana	<i>Bambusa tessallata</i>	Japan
<i>Opuntiaspis</i>	<i>carinata</i>	(Cockerell)	<i>Citrus aurantiifolia</i>	Mexico
	<i>philococcus</i>	Cockerell	<i>Cereus</i> sp.	Mexico
<i>Parlatoria</i>	<i>theae</i>	Cockerell	<i>Thea</i> sp.	Japan
<i>Pinnaspis</i>	sp.	NA	<i>Eugenia</i> sp.	Japan
	<i>theae</i>	(Maskell)	<i>Tea</i> sp.	Taiwan
<i>Poliaspis</i>	sp.	NA	<i>Acacia</i> sp.	Australia
<i>Protodiaspis</i>	Sp.	NA	<i>Quercus</i> sp.	Mexico
	sp.	NA	undetermined	Mexico
<i>Protomorgania</i>	<i>koebelei</i>	Dooley and Evans	<i>Pittosporum</i> sp.	Australia
<i>Pseudaonidia</i>	<i>cingulata</i>	Froggatt	<i>Casuarina</i> sp.	Australia
<i>Pseudaulacaspis</i>	<i>cockerelli</i>	(Cooley)	<i>Eugenia</i> sp.	Hawaii
	<i>cockerelli</i>	(Cooley)	<i>Michaelia figo</i>	California
	<i>cockerelli</i>	(Cooley)	undetermined	Sri Lanka
	<i>cockerelli</i>	(Cooley)	<i>Artocarpus</i> sp.	Hong Kong
	<i>cockerelli</i>	(Cooley)	Palmaceae	Australia
	<i>cockerelli</i>	(Cooley)	undetermined	Sri Lanka





	<i>cockerelli</i>	(Cooley)	<i>Eucalyptus</i> sp.	
	<i>cockerelli</i>	(Cooley)	<i>Ptychosperma</i> sp.	Australia
	<i>cockerelli</i>	(Cooley)	Palmaceae	Australia
<i>Selenaspidus</i>	<i>articulata</i>	Say	<i>Citrus</i> sp.	Mexico
<i>Thysanofiorinia</i>	<i>nephelii</i>	Maskell	<i>Dimocarpus longan</i>	China
<i>Unaspis</i>	sp.	NA	undetermined	Mexico

### List of figures

**Figure 1.** *Protomorgania koebelei* Dooley and Evans. 2012. Illustration by Dr. Greg Evans and image by John Dooley. Original condition of specimens in the collection (left), same specimens slide mounted and illustrated (right).



**Figure 2.** *Gadigaleyrodes froggatti* Dooley and Gillespie 2013. Image by John Dooley







**Figure 3.** Male (left) and female (right) of *Habralictus insularis* Smith-Pardo, 2009.

